

VANILREP4 POLYPEPTIDES AND VANILREP4 POLYNUCLEOTIDES**Cross Referenced Applications**

5 This application is a continuation of co-pending application serial numbers: U.S. Application No 10/395,800, filed 24 March, 2003; U.S. Application No.: 10/286,733, filed 01 November 2002; U.S. Application No.: 10/160,320, filed 30 May 2002, U.S. Application No. 10/039,616, filed 4 January, 2002; U.S. Application No. 09/523,860, filed 13 March 2000.

Field of the Invention

10 This invention relates to newly identified polypeptides and polynucleotides encoding such polypeptides, to their use in diagnosis and in identifying compounds that may be agonists, antagonists that are potentially useful in therapy, and to production of such
15 polypeptides and polynucleotides.

Background of the Invention

20 The drug discovery process is currently undergoing a fundamental revolution as it embraces "functional genomics", that is, high throughput genome- or gene-based biology. This approach as a means to identify genes and gene products as therapeutic targets is rapidly superceding earlier approaches based on "positional cloning". A phenotype, that is a biological function or genetic disease, would be identified and this would then be tracked back to the responsible gene, based on its genetic map position.

25 Functional genomics relies heavily on high-throughput DNA sequencing technologies and the various tools of bioinformatics to identify gene sequences of potential interest from the many molecular biology databases now available. There is a continuing need to identify and characterise further genes and their related polypeptides/proteins, as targets for drug discovery.

Summary of the Invention

30 The present invention relates to VANILREP4, in particular VANILREP4 polypeptides and VANILREP4 polynucleotides, recombinant materials and methods for their production. Such polypeptides and polynucleotides are of interest in relation to methods of treatment of certain diseases, including, but not limited to, pain, chronic pain, neuropathic pain, postoperative pain, rheumatoid arthritic pain, neuralgia, neuropathies, algesia, nerve injury,
35 ischaemia, neurodegeneration, stroke, incontinence, inflammatory disorders, irritable bowel

syndrome, diabetes or obesity, hereinafter referred to as "diseases of the invention". In a further aspect, the invention relates to methods for identifying agonists and antagonists (*e.g.*, inhibitors) using the materials provided by the invention, and treating conditions associated with VANILREP4 imbalance with the identified compounds. In a still further aspect, the invention relates to diagnostic assays for detecting diseases associated with inappropriate VANILREP4 activity or levels.

Description of the Invention

In a first aspect, the present invention relates to VANILREP4 polypeptides. Such polypeptides include:

- (a) an isolated polypeptide encoded by a polynucleotide comprising the sequence of SEQ ID NO:1;
- (b) an isolated polypeptide comprising a polypeptide sequence having at least 95%, 96%, 97%, 98%, or 99% identity to the polypeptide sequence of SEQ ID NO:2;
- (c) an isolated polypeptide comprising the polypeptide sequence of SEQ ID NO:2;
- (d) an isolated polypeptide having at least 95%, 96%, 97%, 98%, or 99% identity to the polypeptide sequence of SEQ ID NO:2;
- (e) the polypeptide sequence of SEQ ID NO:2; and
- (f) an isolated polypeptide having or comprising a polypeptide sequence that has an Identity Index of 0.95, 0.96, 0.97, 0.98, or 0.99 compared to the polypeptide sequence of SEQ ID NO:2;
- (g) fragments and variants of such polypeptides in (a) to (f).

Polypeptides of the present invention are believed to be members of the ion channel family of polypeptides. They are therefore of interest because they are related to the VR1 channel which is associated with the mechanism of action of capsaicin (a vanilloid compound), a constituent of chilli peppers. Capsaicin elicits a sensation of burning pain by selectively activating sensory neurons that convey information about noxious stimuli to the ce.

The biological properties of the VANILREP4 are hereinafter referred to as "biological activity of VANILREP4" or VANILREP4 activity". Preferably, a polypeptide of the present invention exhibits at least one biological activity of VANILREP4. Polypeptides of the present invention also includes variants of the aforementioned polypeptides, including all allelic forms and splice variants. Such polypeptides vary from the reference polypeptide by insertions, deletions, and substitutions that may be conservative or non-conservative, or any combination thereof. Particularly preferred variants are those in which several, for instance from

50 to 30, from 30 to 20, from 20 to 10, from 10 to 5, from 5 to 3, from 3 to 2, from 2 to 1 or 1 amino acids are inserted, substituted, or deleted, in any combination.

Preferred fragments of polypeptides of the present invention include an isolated polypeptide comprising an amino acid sequence having at least 30, 50 or 100 contiguous amino acids from the amino acid sequence of SEQ ID NO: 2, or an isolated polypeptide comprising an amino acid sequence having at least 30, 50 or 100 contiguous amino acids truncated or deleted from the amino acid sequence of SEQ ID NO: 2. Preferred fragments are biologically active fragments that mediate the biological activity of VANILREP4, including those with a similar activity or an improved activity, or with a decreased undesirable activity. Also preferred are those fragments that are antigenic or immunogenic in an animal, especially in a human.

Fragments of the polypeptides of the invention may be employed for producing the corresponding full-length polypeptide by peptide synthesis; therefore, these variants may be employed as intermediates for producing the full-length polypeptides of the invention. The polypeptides of the present invention may be in the form of the "mature" protein or may be a part of a larger protein such as a precursor or a fusion protein. It is often advantageous to include an additional amino acid sequence that contains secretory or leader sequences, pro-sequences, sequences that aid in purification, for instance multiple histidine residues, or an additional sequence for stability during recombinant production.

Polypeptides of the present invention can be prepared in any suitable manner, for instance by isolation from naturally occurring sources, from genetically engineered host cells comprising expression systems (*vide infra*) or by chemical synthesis, using for instance automated peptide synthesizers, or a combination of such methods. Means for preparing such polypeptides are well understood in the art.

In a further aspect, the present invention relates to VANILREP4 polynucleotides. Such polynucleotides include:

- (a) an isolated polynucleotide comprising a polynucleotide sequence having at least 95%, 96%, 97%, 98%, or 99% identity to the polynucleotide sequence of SEQ ID NO:1;
- (b) an isolated polynucleotide comprising the polynucleotide of SEQ ID NO:1;
- (c) an isolated polynucleotide having at least 95%, 96%, 97%, 98%, or 99% identity to the polynucleotide of SEQ ID NO:1;
- (d) the isolated polynucleotide of SEQ ID NO:1;
- (e) an isolated polynucleotide comprising a polynucleotide sequence encoding a polypeptide sequence having at least 95%, 96%, 97%, 98%, or 99% identity to the polypeptide sequence of SEQ ID NO:2;

- (f) an isolated polynucleotide comprising a polynucleotide sequence encoding the polypeptide of SEQ ID NO:2;
- (g) an isolated polynucleotide having a polynucleotide sequence encoding a polypeptide sequence having at least 95%, 96%, 97%, 98%, or 99% identity to the polypeptide sequence of SEQ ID NO:2;
- (h) an isolated polynucleotide encoding the polypeptide of SEQ ID NO:2;
- (i) an isolated polynucleotide having or comprising a polynucleotide sequence that has an Identity Index of 0.95, 0.96, 0.97, 0.98, or 0.99 compared to the polynucleotide sequence of SEQ ID NO:1;
- (j) an isolated polynucleotide having or comprising a polynucleotide sequence encoding a polypeptide sequence that has an Identity Index of 0.95, 0.96, 0.97, 0.98, or 0.99 compared to the polypeptide sequence of SEQ ID NO:2; and polynucleotides that are fragments and variants of the above mentioned polynucleotides or that are complementary to above mentioned polynucleotides, over the entire length thereof.
- Preferred fragments of polynucleotides of the present invention include an isolated polynucleotide comprising an nucleotide sequence having at least 15, 30, 50 or 100 contiguous nucleotides from the sequence of SEQ ID NO: 1, or an isolated polynucleotide comprising an sequence having at least 30, 50 or 100 contiguous nucleotides truncated or deleted from the sequence of SEQ ID NO: 1.
- Preferred variants of polynucleotides of the present invention include splice variants, allelic variants, and polymorphisms, including polynucleotides having one or more single nucleotide polymorphisms (SNPs).
- Polynucleotides of the present invention also include polynucleotides encoding polypeptide variants that comprise the amino acid sequence of SEQ ID NO:2 and in which several, for instance from 50 to 30, from 30 to 20, from 20 to 10, from 10 to 5, from 5 to 3, from 3 to 2, from 2 to 1 or 1 amino acid residues are substituted, deleted or added, in any combination.
- In a further aspect, the present invention provides polynucleotides that are RNA transcripts of the DNA sequences of the present invention. Accordingly, there is provided an RNA polynucleotide that:
- (a) comprises an RNA transcript of the DNA sequence encoding the polypeptide of SEQ ID NO:2;
- (b) is the RNA transcript of the DNA sequence encoding the polypeptide of SEQ ID NO:2;
- (c) comprises an RNA transcript of the DNA sequence of SEQ ID NO:1; or

(d) is the RNA transcript of the DNA sequence of SEQ ID NO:1; and RNA polynucleotides that are complementary thereto.

The polynucleotide sequence of SEQ ID NO:1 shows homology with rat vanilloid receptor, VR1 (M. J. Caterina et al., Nature 389: 816-824, 1997). The polynucleotide sequence of SEQ ID NO:1 is a cDNA sequence that encodes the polypeptide of SEQ ID NO:2. The polynucleotide sequence encoding the polypeptide of SEQ ID NO:2 may be identical to the polypeptide encoding sequence of SEQ ID NO:1 or it may be a sequence other than SEQ ID NO:1, which, as a result of the redundancy (degeneracy) of the genetic code, also encodes the polypeptide of SEQ ID NO:2. The polypeptide of the SEQ ID NO:2 is related to other proteins of the ion channel family, having homology and/or structural similarity with with rat vanilloid receptor, VR1 (M. J. Caterina et al., Nature 389: 816-824, 1997). The nucleotide sequence of SEQ ID NO:4 is a cDNA sequence and comprises a polypeptide encoding sequence (nucleotide 90 to 2705, Exon1 (< 1-58), Exon 2 (59-475), Exon 3 (476-648), Exon 4 (649-801), Exon 5 (802-942), Exon 6 (943-1241), Exon 7 (1242-1421), Exon 8 (1422-1580), Exon 9 (1581-1673), Exon 10 (1674-1747), Exon 11 (1748-1913), Exon 12 (1914-1980), Exon 13 (1981-2297), Exon 14 (2298-2425), Exon 15 (2426-2546), Exon 16 (2547 - >3237)) encoding a polypeptide of 871 amino acids, the polypeptide of SEQ ID NO:2. Knowledge of the intron-exon structure of VANILREP4 can be used for mutation screening, for example as a diagnostic test for diseases which may be caused by alterations of VANILREP4 . The screening of genomic DNA is desirable for the analysis of non-coding regions, such as upstream regulatory regions and intron splice sites. It is also useful in cases where mRNA is not readily available for mutation analysis. Knowledge of the genomic structure is also important for the generation of animal models. Such models may be used to study the function of VANILREP4 and for drug screening studies. For example, mouse knock-out models typically have a selection marker, which upon insertion into a coding exon, ablate the functioning of the targeted allele. The genomic structure may also be useful in analysing possible splice variants of VANILREP4. Splice variants are important because they may have different functions and different expression patterns.

Preferred polypeptides and polynucleotides of the present invention are expected to have, *inter alia*, similar biological functions/properties to their homologous polypeptides and polynucleotides. Furthermore, preferred polypeptides and polynucleotides of the present invention have at least one VANILREP4 activity.

The present invention also relates to partial or other polynucleotide and polypeptide sequences which were first identified prior to the determination of the corresponding full length sequence of SEQ ID NO:4. Accordingly, in a further aspect, the present invention provides for an isolated polynucleotide which:

(a) comprises a nucleotide sequence which has at least 95% identity, preferably at least 97-99% identity to SEQ ID NO:3 over the entire length of SEQ ID NO:3;

(b) has a nucleotide sequence which has at least 95% identity, preferably at least 97-99% identity, to SEQ ID NO:3 over the entire length of SEQ ID NO:3; or

5 (c) the polynucleotide of SEQ ID NO:3.

The nucleotide sequence of SEQ ID NO:3 is derived from a single cDNA clone. The 5' 11bp of SEQ ID NO:3 are at variance with the equivalent region of SEQ ID NO:4, and most likely result from nucleotide sequence reading errors of this first cDNA clone, or are the result of a cloning artefact. It is well known in the art that cDNA cloning and sequencing can be vulnerable to such problems (see Aaronson, J.S. et al. *Genome Research* 6:829-845 1996).
 10 The entire coding regions of the polynucleotides of SEQ ID NO:1, SEQ ID NO:3 and SEQ ID NO:4 are identical, and hence the polypeptides encoded therefrom are also identical, having the amino acid sequence of SEQ ID NO:2.

Polynucleotides of the present invention may be obtained using standard cloning and screening techniques from a cDNA library derived from mRNA in cells of human brain, germ cell, whole embryo, heart, kidney, prostate, lung, uterus, glioblastoma, adenocarcinoma, senescent fibroblast and osteoarthritic cartilage, (see for instance, Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989)). Polynucleotides of the invention can also be obtained from natural
 15 sources such as genomic DNA libraries or can be synthesized using well known and commercially available techniques.

When polynucleotides of the present invention are used for the recombinant production of polypeptides of the present invention, the polynucleotide may include the coding sequence for the mature polypeptide, by itself, or the coding sequence for the mature
 25 polypeptide in reading frame with other coding sequences, such as those encoding a leader or secretory sequence, a pre-, or pro- or prepro- protein sequence, or other fusion peptide portions. For example, a marker sequence that facilitates purification of the fused polypeptide can be encoded. In certain preferred embodiments of this aspect of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz
 30 *et al.*, *Proc Natl Acad Sci USA* (1989) 86:821-824, or is an HA tag. The polynucleotide may also contain non-coding 5' and 3' sequences, such as transcribed, non-translated sequences, splicing and polyadenylation signals, ribosome binding sites and sequences that stabilize mRNA.

Polynucleotides that are identical, or have sufficient identity to a polynucleotide
 35 sequence of SEQ ID NO:1, may be used as hybridization probes for cDNA and genomic DNA

or as primers for a nucleic acid amplification reaction (for instance, PCR). Such probes and primers may be used to isolate full-length cDNAs and genomic clones encoding polypeptides of the present invention and to isolate cDNA and genomic clones of other genes (including genes encoding paralogs from human sources and orthologs and paralogs from species other than human) that have a high sequence similarity to SEQ ID NO:1, typically at least 95% identity. Preferred probes and primers will generally comprise at least 15 nucleotides, preferably, at least 30 nucleotides and may have at least 50, if not at least 100 nucleotides. Particularly preferred probes will have between 30 and 50 nucleotides. Particularly preferred primers will have between 20 and 25 nucleotides.

A polynucleotide encoding a polypeptide of the present invention, including homologs from species other than human, may be obtained by a process comprising the steps of screening a library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO: 1 or a fragment thereof, preferably of at least 15 nucleotides; and isolating full-length cDNA and genomic clones containing said polynucleotide sequence. Such hybridization techniques are well known to the skilled artisan. Preferred stringent hybridization conditions include overnight incubation at 42°C in a solution comprising: 50% formamide, 5xSSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10 % dextran sulfate, and 20 microgram/ml denatured, sheared salmon sperm DNA; followed by washing the filters in 0.1x SSC at about 65°C. Thus the present invention also includes isolated polynucleotides, preferably with a nucleotide sequence of at least 100, obtained by screening a library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO:1 or a fragment thereof, preferably of at least 15 nucleotides.

The skilled artisan will appreciate that, in many cases, an isolated cDNA sequence will be incomplete, in that the region coding for the polypeptide does not extend all the way through to the 5' terminus. This is a consequence of reverse transcriptase, an enzyme with inherently low "processivity" (a measure of the ability of the enzyme to remain attached to the template during the polymerisation reaction), failing to complete a DNA copy of the mRNA template during first strand cDNA synthesis.

There are several methods available and well known to those skilled in the art to obtain full-length cDNAs, or extend short cDNAs, for example those based on the method of Rapid Amplification of cDNA ends (RACE) (see, for example, Frohman et al., Proc Nat Acad Sci USA 85, 8998-9002, 1988). Recent modifications of the technique, exemplified by the Marathon (trade mark) technology (Clontech Laboratories Inc.) for example, have significantly simplified the search for longer cDNAs. In the Marathon (trade mark)

technology, cDNAs have been prepared from mRNA extracted from a chosen tissue and an 'adaptor' sequence ligated onto each end. Nucleic acid amplification (PCR) is then carried out to amplify the "missing" 5' end of the cDNA using a combination of gene specific and adaptor specific oligonucleotide primers. The PCR reaction is then repeated using 'nested' primers, that is, primers designed to anneal within the amplified product (typically an adaptor specific primer that anneals further 3' in the adaptor sequence and a gene specific primer that anneals further 5' in the known gene sequence). The products of this reaction can then be analysed by DNA sequencing and a full-length cDNA constructed either by joining the product directly to the existing cDNA to give a complete sequence, or carrying out a separate full-length PCR using the new sequence information for the design of the 5' primer.

Recombinant polypeptides of the present invention may be prepared by processes well known in the art from genetically engineered host cells comprising expression systems. Accordingly, in a further aspect, the present invention relates to expression systems comprising a polynucleotide or polynucleotides of the present invention, to host cells which are genetically engineered with such expression systems and to the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention.

For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof for polynucleotides of the present invention.

Polynucleotides may be introduced into host cells by methods described in many standard laboratory manuals, such as Davis et al., Basic Methods in Molecular Biology (1986) and Sambrook *et al.*(*ibid*). Preferred methods of introducing polynucleotides into host cells include, for instance, calcium phosphate transfection, DEAE-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction or infection.

Representative examples of appropriate hosts include bacterial cells, such as *Streptococci*, *Staphylococci*, *E. coli*, *Streptomyces* and *Bacillus subtilis* cells; fungal cells, such as yeast cells and *Aspergillus* cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, HEK 293 and Bowes melanoma cells; and plant cells.

A great variety of expression systems can be used, for instance, chromosomal, episomal and virus-derived systems, *e.g.*, vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from

combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression systems may contain control regions that regulate as well as engender expression. Generally, any system or vector that is able to maintain, propagate or express a polynucleotide to produce a polypeptide in a host may be used.

5 The appropriate polynucleotide sequence may be inserted into an expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook *et al.*, (*ibid*). Appropriate secretion signals may be incorporated into the desired polypeptide to allow secretion of the translated protein into the lumen of the endoplasmic reticulum, the periplasmic space or the extracellular environment. These signals may be endogenous to the
10 polypeptide or they may be heterologous signals.

If a polypeptide of the present invention is to be expressed for use in screening assays, it is generally preferred that the polypeptide be produced at the surface of the cell. In this event, the cells may be harvested prior to use in the screening assay. If the polypeptide is secreted into the medium, the medium can be recovered in order to recover and purify the
15 polypeptide. If produced intracellularly, the cells must first be lysed before the polypeptide is recovered.

Polypeptides of the present invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography,
20 hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well known techniques for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during intracellular synthesis, isolation and/or purification.

25 Polynucleotides of the present invention may be used as diagnostic reagents, through detecting mutations in the associated gene. Detection of a mutated form of the gene characterised by the polynucleotide of SEQ ID NO:1 in the cDNA or genomic sequence and which is associated with a dysfunction will provide a diagnostic tool that can add to, or define, a diagnosis of a disease, or susceptibility to a disease, which results from under-expression, over-
30 expression or altered spatial or temporal expression of the gene. Individuals carrying mutations in the gene may be detected at the DNA level by a variety of techniques well known in the art.

Nucleic acids for diagnosis may be obtained from a subject's cells, such as from blood, urine, saliva, tissue biopsy or autopsy material. The genomic DNA may be used directly for detection or it may be amplified enzymatically by using PCR, preferably RT-PCR, or other
35 amplification techniques prior to analysis. RNA or cDNA may also be used in similar fashion.

Deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to labeled VANILREP4 nucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting
 5 temperatures. DNA sequence difference may also be detected by alterations in the electrophoretic mobility of DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing (see, for instance, Myers *et al.*, Science (1985) 230:1242). Sequence changes at specific locations may also be revealed by nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method (see Cotton *et al.*, Proc Natl Acad Sci USA
 10 (1985) 85: 4397-4401).

An array of oligonucleotides probes comprising VANILREP4 polynucleotide sequence or fragments thereof can be constructed to conduct efficient screening of *e.g.*, genetic mutations. Such arrays are preferably high density arrays or grids. Array technology methods are well known and have general applicability and can be used to address a variety of questions in
 15 molecular genetics including gene expression, genetic linkage, and genetic variability, see, for example, M.Chee et al., Science, 274, 610-613 (1996) and other references cited therein.

Detection of abnormally decreased or increased levels of polypeptide or mRNA expression may also be used for diagnosing or determining susceptibility of a subject to a disease of the invention. Decreased or increased expression can be measured at the RNA
 20 level using any of the methods well known in the art for the quantitation of polynucleotides, such as, for example, nucleic acid amplification, for instance PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods. Assay techniques that can be used to determine levels of a protein, such as a polypeptide of the present invention, in a sample derived from a host are well-known to those of skill in the art. Such assay methods include
 25 radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

Thus in another aspect, the present invention relates to a diagnostic kit comprising:
 (a) a polynucleotide of the present invention, preferably the nucleotide sequence of SEQ ID NO: 1, or a fragment or an RNA transcript thereof;
 (b) a nucleotide sequence complementary to that of (a);
 30 (c) a polypeptide of the present invention, preferably the polypeptide of SEQ ID NO:2 or a fragment thereof; or
 (d) an antibody to a polypeptide of the present invention, preferably to the polypeptide of SEQ ID NO:2.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component. Such a kit will be of use in diagnosing a disease or susceptibility to a disease, particularly diseases of the invention, amongst others.

The polynucleotide sequences of the present invention are valuable for chromosome localisation studies. The sequence is specifically targeted to, and can hybridize with, a particular location on an individual human chromosome. The mapping of relevant sequences to chromosomes according to the present invention is an important first step in correlating those sequences with gene associated disease. Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found in, for example, V. McKusick, Mendelian Inheritance in Man (available on-line through Johns Hopkins University Welch Medical Library). The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (co-inheritance of physically adjacent genes). Precise human chromosomal localisations for a genomic sequence (gene fragment etc.) can be determined using Radiation Hybrid (RH) Mapping (Walter, M. Spillett, D., Thomas, P., Weissenbach, J., and Goodfellow, P., (1994) A method for constructing radiation hybrid maps of whole genomes, *Nature Genetics* 7, 22-28). A number of RH panels are available from Research Genetics (Huntsville, AL, USA) e.g. the GeneBridge4 RH panel (*Hum Mol Genet* 1996 Mar;5(3):339-46 A radiation hybrid map of the human genome. Gyapay G, Schmitt K, Fizames C, Jones H, Vega-Czarny N, Spillett D, Muselet D, Prud'Homme JF, Dib C, Auffray C, Morissette J, Weissenbach J, Goodfellow PN). To determine the chromosomal location of a gene using this panel, 93 PCRs are performed using primers designed from the gene of interest on RH DNAs. Each of these DNAs contains random human genomic fragments maintained in a hamster background (human / hamster hybrid cell lines). These PCRs result in 93 scores indicating the presence or absence of the PCR product of the gene of interest. These scores are compared with scores created using PCR products from genomic sequences of known location. This comparison is conducted at <http://www.genome.wi.mit.edu/>. The gene of the present invention maps to human chromosome 12q24.1.

The polynucleotide sequences of the present invention are also valuable tools for tissue expression studies. Such studies allow the determination of expression patterns of polynucleotides of the present invention which may give an indication as to the expression patterns of the encoded polypeptides in tissues, by detecting the mRNAs that encode them. The techniques used are well known in the art and include in situ hybridisation techniques to clones arrayed on a grid, such as cDNA microarray hybridisation (Schena *et al*, *Science*, 270, 467-470,

1995 and Shalon *et al*, Genome Res, 6, 639-645, 1996) and nucleotide amplification techniques such as PCR. A preferred method uses the TAQMAN (Trade mark) technology available from Perkin Elmer. Results from these studies can provide an indication of the normal function of the polypeptide in the organism. In addition, comparative studies of the normal expression pattern of mRNAs with that of mRNAs encoded by an alternative form of the same gene (for example, one having an alteration in polypeptide coding potential or a regulatory mutation) can provide valuable insights into the role of the polypeptides of the present invention, or that of inappropriate expression thereof in disease. Such inappropriate expression may be of a temporal, spatial or simply quantitative nature.

The polynucleotides of the present invention are expressed in several tissues, including human brain, germ cell, whole embryo, heart, kidney, prostate, lung, uterus, glioblastoma, adenocarcinoma, senescent fibroblast, osteoarthritic cartilage and sensory ganglia.

A further aspect of the present invention relates to antibodies. The polypeptides of the invention or their fragments, or cells expressing them, can be used as immunogens to produce antibodies that are immunospecific for polypeptides of the present invention. The term "immunospecific" means that the antibodies have substantially greater affinity for the polypeptides of the invention than their affinity for other related polypeptides in the prior art.

Antibodies generated against polypeptides of the present invention may be obtained by administering the polypeptides or epitope-bearing fragments, or cells to an animal, preferably a non-human animal, using routine protocols. For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler, G. and Milstein, C., Nature (1975) 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor *et al.*, Immunology Today (1983) 4:72) and the EBV-hybridoma technique (Cole *et al.*, Monoclonal Antibodies and Cancer Therapy, 77-96, Alan R. Liss, Inc., 1985).

Techniques for the production of single chain antibodies, such as those described in U.S. Patent No. 4,946,778, can also be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms, including other mammals, may be used to express humanized antibodies.

The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptide or to purify the polypeptides by affinity chromatography. Antibodies against polypeptides of the present invention may also be employed to treat diseases of the invention, amongst others.

Polypeptides and polynucleotides of the present invention may also be used as vaccines. Accordingly, in a further aspect, the present invention relates to a method for

inducing an immunological response in a mammal that comprises inoculating the mammal with a polypeptide of the present invention, adequate to produce antibody and/or T cell immune response, including, for example, cytokine-producing T cells or cytotoxic T cells, to protect said animal from disease, whether that disease is already established within the individual or not. An immunological response in a mammal may also be induced by a method comprises delivering a polypeptide of the present invention *via* a vector directing expression of the polynucleotide and coding for the polypeptide *in vivo* in order to induce such an immunological response to produce antibody to protect said animal from diseases of the invention. One way of administering the vector is by accelerating it into the desired cells as a coating on particles or otherwise. Such nucleic acid vector may comprise DNA, RNA, a modified nucleic acid, or a DNA/RNA hybrid. For use a vaccine, a polypeptide or a nucleic acid vector will be normally provided as a vaccine formulation (composition). The formulation may further comprise a suitable carrier. Since a polypeptide may be broken down in the stomach, it is preferably administered parenterally (for instance, subcutaneous, intramuscular, intravenous, or intradermal injection). Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions that may contain anti-oxidants, buffers, bacteriostats and solutes that render the formulation isotonic with the blood of the recipient; and aqueous and non-aqueous sterile suspensions that may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

Polypeptides of the present invention have one or more biological functions that are of relevance in one or more disease states, in particular the diseases of the invention hereinbefore mentioned. It is therefore useful to to identify compounds that stimulate or inhibit the function or level of the polypeptide. Accordingly, in a further aspect, the present invention provides for a method of screening compounds to identify those that stimulate or inhibit the function or level of the polypeptide. Such methods identify agonists or antagonists that may be employed for therapeutic and prophylactic purposes for such diseases of the invention as hereinbefore mentioned. Compounds may be identified from a variety of sources, for example, cells, cell-free preparations, chemical libraries, collections of chemical compounds, and natural product mixtures. Such agonists or antagonists so-identified may be natural or modified substrates,

ligands, receptors, enzymes, etc., as the case may be, of the polypeptide; a structural or functional mimetic thereof (see Coligan *et al.*, Current Protocols in Immunology 1(2):Chapter 5 (1991)) or a small molecule. Such small molecules preferably have a molecular weight below 2,000 daltons, more preferably between 300 and 1,000 daltons, and most preferably between 400 and 700 daltons. It is preferred that these small molecules are organic molecules.

The screening method may simply measure the binding of a candidate compound to the polypeptide, or to cells or membranes bearing the polypeptide, or a fusion protein thereof, by means of a label directly or indirectly associated with the candidate compound. Alternatively, the screening method may involve measuring or detecting (qualitatively or quantitatively) the competitive binding of a candidate compound to the polypeptide against a labeled competitor (*e.g.* agonist or antagonist). Further, these screening methods may test whether the candidate compound results in a signal generated by activation or inhibition of the polypeptide, using detection systems appropriate to the cells bearing the polypeptide. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed. Further, the screening methods may simply comprise the steps of mixing a candidate compound with a solution containing a polypeptide of the present invention, to form a mixture, measuring a VANILREP4 activity in the mixture, and comparing the VANILREP4 activity of the mixture to a control mixture which contains no candidate compound.

Polypeptides of the present invention may be employed in conventional low capacity screening methods and also in high-throughput screening (HTS) formats. Such HTS formats include not only the well-established use of 96- and, more recently, 384-well micotiter plates but also emerging methods such as the nanowell method described by Schullek *et al.*, Anal Biochem., 246, 20-29, (1997).

Fusion proteins, such as those made from Fc portion and VANILREP4 polypeptide, as hereinbefore described, can also be used for high-throughput screening assays to identify antagonists for the polypeptide of the present invention (see D. Bennett *et al.*, J Mol Recognition, 8:52-58 (1995); and K. Johanson *et al.*, J Biol Chem, 270(16):9459-9471 (1995)).

The polynucleotides, polypeptides and antibodies to the polypeptide of the present invention may also be used to configure screening methods for detecting the effect of added compounds on the production of mRNA and polypeptide in cells. For example, an ELISA assay may be constructed for measuring secreted or cell associated levels of polypeptide using monoclonal and polyclonal antibodies by standard methods known in the art. This can

be used to discover agents that may inhibit or enhance the production of polypeptide (also called antagonist or agonist, respectively) from suitably manipulated cells or tissues.

A polypeptide of the present invention may be used to identify membrane bound or soluble receptors, if any, through standard receptor binding techniques known in the art.

- 5 These include, but are not limited to, ligand binding and crosslinking assays in which the polypeptide is labeled with a radioactive isotope (for instance, ^{125}I), chemically modified (for instance, biotinylated), or fused to a peptide sequence suitable for detection or purification, and incubated with a source of the putative receptor (cells, cell membranes, cell supernatants, tissue extracts, bodily fluids). Other methods include biophysical techniques
10 such as surface plasmon resonance and spectroscopy. These screening methods may also be used to identify agonists and antagonists of the polypeptide that compete with the binding of the polypeptide to its receptors, if any. Standard methods for conducting such assays are well understood in the art.

- Examples of antagonists of polypeptides of the present invention include antibodies or,
15 in some cases, oligonucleotides or proteins that are closely related to the ligands, substrates, receptors, enzymes, etc., as the case may be, of the polypeptide, *e.g.*, a fragment of the ligands, substrates, receptors, enzymes, etc.; or a small molecule that bind to the polypeptide of the present invention but do not elicit a response, so that the activity of the polypeptide is prevented.

- Screening methods may also involve the use of transgenic technology and
20 VANILREP4 gene. The art of constructing transgenic animals is well established. For example, the VANILREP4 gene may be introduced through microinjection into the male pronucleus of fertilized oocytes, retroviral transfer into pre- or post-implantation embryos, or injection of genetically modified, such as by electroporation, embryonic stem cells into host blastocysts. Particularly useful transgenic animals are so-called "knock-in" animals in which
25 an animal gene is replaced by the human equivalent within the genome of that animal. Knock-in transgenic animals are useful in the drug discovery process, for target validation, where the compound is specific for the human target. Other useful transgenic animals are so-called "knock-out" animals in which the expression of the animal ortholog of a polypeptide of the present invention and encoded by an endogenous DNA sequence in a cell is partially
30 or completely annulled. The gene knock-out may be targeted to specific cells or tissues, may occur only in certain cells or tissues as a consequence of the limitations of the technology, or may occur in all, or substantially all, cells in the animal. Transgenic animal technology also offers a whole animal expression-cloning system in which introduced genes are expressed to give large amounts of polypeptides of the present invention

Screening kits for use in the above described methods form a further aspect of the present invention. Such screening kits comprise:

- (a) a polypeptide of the present invention;
 - (b) a recombinant cell expressing a polypeptide of the present invention;
 - 5 (c) a cell membrane expressing a polypeptide of the present invention; or
 - (d) an antibody to a polypeptide of the present invention;
- which polypeptide is preferably that of SEQ ID NO:2.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

10

Glossary

The following definitions are provided to facilitate understanding of certain terms used frequently hereinbefore.

- “Antibodies” as used herein includes polyclonal and monoclonal antibodies, chimeric, single chain, and humanized antibodies, as well as Fab fragments, including the products of an Fab or other immunoglobulin expression library.

- “Isolated” means altered “by the hand of man” from its natural state, *i.e.*, if it occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living organism is not “isolated,” but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is “isolated”, as the term is employed herein. Moreover, a polynucleotide or polypeptide that is introduced into an organism by transformation, genetic manipulation or by any other recombinant method is “isolated” even if it is still present in said organism, which organism may be living or non-living.

- “Polynucleotide” generally refers to any polyribonucleotide (RNA) or polydeoxribonucleotide (DNA), which may be unmodified or modified RNA or DNA. “Polynucleotides” include, without limitation, single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, “polynucleotide” refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The term “polynucleotide” also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. “Modified” bases include, for

example, tritylated bases and unusual bases such as inosine. A variety of modifications may be made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells. "Polynucleotide" also
 5 embraces relatively short polynucleotides, often referred to as oligonucleotides.

"Polypeptide" refers to any polypeptide comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres.

"Polypeptide" refers to both short chains, commonly referred to as peptides, oligopeptides or oligomers, and to longer chains, generally referred to as proteins. Polypeptides may contain
 10 amino acids other than the 20 gene-encoded amino acids. "Polypeptides" include amino acid sequences modified either by natural processes, such as post-translational processing, or by chemical modification techniques that are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications may occur anywhere in a polypeptide, including the
 15 peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present to the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be branched as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched and branched cyclic polypeptides may
 20 result from post-translation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, biotinylation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond
 25 formation, demethylation, formation of covalent cross-links, formation of cystine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination (see, for instance,
 30 *Proteins - Structure and Molecular Properties*, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York, 1993; Wold, F., *Post-translational Protein Modifications: Perspectives and Prospects*, 1-12, in *Post-translational Covalent Modification of Proteins*, B. C. Johnson, Ed., Academic Press, New York, 1983; Seifter *et al.*, "Analysis for protein modifications and nonprotein cofactors", *Meth Enzymol*, 182, 626-646, 1990, and Rattan *et al.*, "Protein

Synthesis: Post-translational Modifications and Aging", Ann NY Acad Sci, 663, 48-62, 1992).

"Fragment" of a polypeptide sequence refers to a polypeptide sequence that is shorter than the reference sequence but that retains essentially the same biological function or activity as the reference polypeptide. "Fragment" of a polynucleotide sequence refers to a polynucleotide sequence that is shorter than the reference sequence of SEQ ID NO:1.

"Variant" refers to a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide, but retains the essential properties thereof. A typical variant of a polynucleotide differs in nucleotide sequence from the reference polynucleotide.

Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from the reference polypeptide. Generally, alterations are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, insertions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. Typical conservative substitutions include Gly, Ala; Val, Ile, Leu; Asp, Glu; Asn, Gln; Ser, Thr; Lys, Arg; and Phe and Tyr. A variant of a polynucleotide or polypeptide may be naturally occurring such as an allele, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques or by direct synthesis. Also included as variants are polypeptides having one or more post-translational modifications, for instance glycosylation, phosphorylation, methylation, ADP ribosylation and the like. Embodiments include methylation of the N-terminal amino acid, phosphorylations of serines and threonines and modification of C-terminal glycines.

"Allele" refers to one of two or more alternative forms of a gene occurring at a given locus in the genome.

"Polymorphism" refers to a variation in nucleotide sequence (and encoded polypeptide sequence, if relevant) at a given position in the genome within a population.

"Single Nucleotide Polymorphism" (SNP) refers to the occurrence of nucleotide variability at a single nucleotide position in the genome, within a population. An SNP may occur within a gene or within intergenic regions of the genome. SNPs can be assayed using Allele Specific Amplification (ASA). For the process at least 3 primers are required. A

common primer is used in reverse complement to the polymorphism being assayed. This common primer can be between 50 and 1500 bps from the polymorphic base. The other two (or more) primers are identical to each other except that the final 3' base wobbles to match one of the two (or more) alleles that make up the polymorphism. Two (or more) PCR reactions are then conducted on sample DNA, each using the common primer and one of the Allele Specific Primers.

"Splice Variant" as used herein refers to cDNA molecules produced from RNA molecules initially transcribed from the same genomic DNA sequence but which have undergone alternative RNA splicing. Alternative RNA splicing occurs when a primary RNA transcript undergoes splicing, generally for the removal of introns, which results in the production of more than one mRNA molecule each of that may encode different amino acid sequences. The term splice variant also refers to the proteins encoded by the above cDNA molecules.

"Identity" reflects a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, determined by comparing the sequences. In general, identity refers to an exact nucleotide to nucleotide or amino acid to amino acid correspondence of the two polynucleotide or two polypeptide sequences, respectively, over the length of the sequences being compared.

"% Identity" - For sequences where there is not an exact correspondence, a "% identity" may be determined. In general, the two sequences to be compared are aligned to give a maximum correlation between the sequences. This may include inserting "gaps" in either one or both sequences, to enhance the degree of alignment. A % identity may be determined over the whole length of each of the sequences being compared (so-called global alignment), that is particularly suitable for sequences of the same or very similar length, or over shorter, defined lengths (so-called local alignment), that is more suitable for sequences of unequal length.

"Similarity" is a further, more sophisticated measure of the relationship between two polypeptide sequences. In general, "similarity" means a comparison between the amino acids of two polypeptide chains, on a residue by residue basis, taking into account not only exact correspondences between a between pairs of residues, one from each of the sequences being compared (as for identity) but also, where there is not an exact correspondence, whether, on an evolutionary basis, one residue is a likely substitute for the other. This likelihood has an associated "score" from which the "% similarity" of the two sequences can then be determined.

Methods for comparing the identity and similarity of two or more sequences are well known in the art. Thus for instance, programs available in the Wisconsin Sequence Analysis Package, version 9.1 (Devereux J et al, Nucleic Acids Res, 12, 387-395, 1984, available from Genetics Computer Group, Madison, Wisconsin, USA), for example the programs BESTFIT and GAP, may be used to determine the % identity between two polynucleotides and the % identity and the % similarity between two polypeptide sequences. BESTFIT uses the "local homology" algorithm of Smith and Waterman (J Mol Biol, 147,195-197, 1981, Advances in Applied Mathematics, 2, 482-489, 1981) and finds the best single region of similarity between two sequences. BESTFIT is more suited to comparing two polynucleotide or two polypeptide sequences that are dissimilar in length, the program assuming that the shorter sequence represents a portion of the longer. In comparison, GAP aligns two sequences, finding a "maximum similarity", according to the algorithm of Needleman and Wunsch (J Mol Biol, 48, 443-453, 1970). GAP is more suited to comparing sequences that are approximately the same length and an alignment is expected over the entire length. Preferably, the parameters "Gap Weight" and "Length Weight" used in each program are 50 and 3, for polynucleotide sequences and 12 and 4 for polypeptide sequences, respectively. Preferably, % identities and similarities are determined when the two sequences being compared are optimally aligned.

Other programs for determining identity and/or similarity between sequences are also known in the art, for instance the BLAST family of programs (Altschul S F et al, J Mol Biol, 215, 403-410, 1990, Altschul S F et al, Nucleic Acids Res., 25:389-3402, 1997, available from the National Center for Biotechnology Information (NCBI), Bethesda, Maryland, USA and accessible through the home page of the NCBI at www.ncbi.nlm.nih.gov) and FASTA (Pearson W R, Methods in Enzymology, 183, 63-99, 1990; Pearson W R and Lipman D J, Proc Nat Acad Sci USA, 85, 2444-2448, 1988, available as part of the Wisconsin Sequence Analysis Package).

Preferably, the BLOSUM62 amino acid substitution matrix (Henikoff S and Henikoff J G, Proc. Nat. Acad Sci. USA, 89, 10915-10919, 1992) is used in polypeptide sequence comparisons including where nucleotide sequences are first translated into amino acid sequences before comparison.

Preferably, the program BESTFIT is used to determine the % identity of a query polynucleotide or a polypeptide sequence with respect to a reference polynucleotide or a polypeptide sequence, the query and the reference sequence being optimally aligned and the parameters of the program set at the default value, as hereinbefore described.

"Identity Index" is a measure of sequence relatedness which may be used to compare a candidate sequence (polynucleotide or polypeptide) and a reference sequence. Thus, for instance, a candidate polynucleotide sequence having, for example, an Identity Index of 0.95 compared to a reference polynucleotide sequence is identical to the reference sequence

5 except that the candidate polynucleotide sequence may include on average up to five differences per each 100 nucleotides of the reference sequence. Such differences are selected from the group consisting of at least one nucleotide deletion, substitution, including transition and transversion, or insertion. These differences may occur at the 5' or 3' terminal positions of the reference polynucleotide sequence or anywhere between these terminal positions,
10 interspersed either individually among the nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence. In other words, to obtain a polynucleotide sequence having an Identity Index of 0.95 compared to a reference polynucleotide sequence, an average of up to 5 in every 100 of the nucleotides of the in the reference sequence may be deleted, substituted or inserted, or any combination thereof, as
15 hereinbefore described. The same applies *mutatis mutandis* for other values of the Identity Index, for instance 0.96, 0.97, 0.98 and 0.99.

Similarly, for a polypeptide, a candidate polypeptide sequence having, for example, an Identity Index of 0.95 compared to a reference polypeptide sequence is identical to the reference sequence except that the polypeptide sequence may include an average of up to five
20 differences per each 100 amino acids of the reference sequence. Such differences are selected from the group consisting of at least one amino acid deletion, substitution, including conservative and non-conservative substitution, or insertion. These differences may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or anywhere between these terminal positions, interspersed either individually among the amino acids in
25 the reference sequence or in one or more contiguous groups within the reference sequence. In other words, to obtain a polypeptide sequence having an Identity Index of 0.95 compared to a reference polypeptide sequence, an average of up to 5 in every 100 of the amino acids in the reference sequence may be deleted, substituted or inserted, or any combination thereof, as
30 hereinbefore described. The same applies *mutatis mutandis* for other values of the Identity Index, for instance 0.96, 0.97, 0.98 and 0.99.

The relationship between the number of nucleotide or amino acid differences and the Identity Index may be expressed in the following equation:

$$n_a \leq x_a - (x_a \cdot I),$$

in which:

35 n_a is the number of nucleotide or amino acid differences,

x_a is the total number of nucleotides or amino acids in SEQ ID NO:1 or SEQ ID NO:2, respectively,

I is the Identity Index ,

• is the symbol for the multiplication operator, and

- 5 in which any non-integer product of x_a and I is rounded down to the nearest integer prior to subtracting it from x_a .

"Homolog" is a generic term used in the art to indicate a polynucleotide or polypeptide sequence possessing a high degree of sequence relatedness to a reference sequence. Such relatedness may be quantified by determining the degree of identity and/or
10 similarity between the two sequences as hereinbefore defined. Falling within this generic term are the terms "ortholog", and "paralog". "Ortholog" refers to a polynucleotide or polypeptide that is the functional equivalent of the polynucleotide or polypeptide in another species. "Paralog" refers to a polynucleotide or polypeptide that within the same species which is functionally similar.

15 "Fusion protein" refers to a protein encoded by two, often unrelated, fused genes or fragments thereof. In one example, EP-A-0 464 533-A discloses fusion proteins comprising various portions of constant region of immunoglobulin molecules together with another human protein or part thereof. In many cases, employing an immunoglobulin Fc region as a part of a fusion protein is advantageous for use in therapy and diagnosis resulting in, for
20 example, improved pharmacokinetic properties [see, *e.g.*, EP-A 0232 262]. On the other hand, for some uses it would be desirable to be able to delete the Fc part after the fusion protein has been expressed, detected and purified.

25 All publications and references, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference in their entirety as if each individual publication or reference were specifically and individually indicated to be incorporated by reference herein as being fully set forth. Any patent application to which this application claims priority is also incorporated by reference herein in its entirety in the manner described above for publications and references.

30

Examples

Example 1: Taqman analysis of mRNA tissue distribution

Expression pattern of VR-4 was investigated using Taqman fluorescent PCR (Perkin Elmer)

- 5 and human cDNAs prepared from various brain areas and peripheral tissues. All Taqman analysis was carried out according to the manufacturers instructions using the following oligonucleotides:

	VR-4 labelled probe:	5'-ATGAGGACCAGACCAACTGCA
10	VR-4 forward primer	5'-GGAGGAAGGTGCTGAAGGTCTC
	VR-4 reverse primer	5'-CACTTACCCCTCGTGCCGTGACAG

- 15 Signals obtained by Taqman PCR were normalised to the housekeeping gene cyclophilin to correct for differences in RNA integrity and quantity. Expression of VR-4 was highest in kidney and generally higher in many peripheral tissues (e.g. liver, pancreas, placenta and prostate) than in CNS. Highest levels of VR-4 in CNS were observed in corpus callosum, hippocampus, spinal cord and pituitary gland.

The tissues investigated were:

- 20 1) amygdala, 2) caudate nucleus, 3) cerebellum, 4) corpus callosum, 5) frontal cortex, 6) occipital cortex, 7) temporal cortex, 8) hippocampus, 9) hypothalamus, 10) nucleus accumbens, 11) putamen, 12) substantia nigra, 13) thalamus, 14) foetal brain, 15) spinal cord, 16) pituitary gland, 17) whole brain, 18) heart, 19) liver, 20) lung, 21) skeletal muscle, 22) kidney, 23) pancreas, 24) spleen, 25) small intestine, 26) placenta, 27) testis, 28) stomach, 29) prostate, 30) uterus

25

Example 2 – Activation of VANILREP4 by phorbol esters

- HEK293 cells transiently expressing hVR1 or hVR4 were seeded into FLIPR96 plates at 25,000cells/well and cultured overnight. The cells were then loaded in medium containing 4microM Fluo-3 for 2hrs, at room temperature, in the dark. The cell plates were then washed 4x with Tyrode containing 1.5mM calcium and 0.2% BSA. Agonist and antagonist plates were prepared in the same buffer. Cells were preincubated with either antagonist or buffer at room temperature for 30 mins. Agonists additions and calcium measurements were made in the FLIPR (Smart *et al.*, (2000) Br J. Pharmacol. 129, 227-230).

- 35 Both phorbol 12-myristate 13-acetate (PMA) and 4 α -phorbol-12,13-didecanoate (4 α PDD) increased intracellular calcium in HEK293-hVR4 cells (Table 1) but were without

effect in wild type HEK293 cells or in cells transfected with empty vector. PMA also activated VR1, but was only a partial agonist (E_{max} 0.46) compared to capsaicin and RTX. 4 α PDD was inactive at VR1 (Table 1).

5 Table 1

	pEC50			
	wild type	empty vector	hVR1	hVR4
RTX	IA	IA	8.93 \pm 0.20	IA
capsaicin	IA	IA	7.48 \pm 0.12	IA
PMA	IA	IA	7.86 \pm 0.06	6.64 \pm 0.06
4 α PDD	IA	IA	IA	5.73 \pm 0.06

Data are mean \pm s.e.mean, where n=3-5. IA= inactive

In conclusion, 4 α PDD acts as a VR4 selective agonist.

SEQUENCE INFORMATION

SEQ ID NO:1

ATGGCGGATTCCAGCGAAGGCCCCCGCGCGGGGCCCGGGAGGTGGCTGAGCTCCCCGGG
 GATGAGAGTGGCACCCAGGTGGGGAGGCTTTTCCTCTCTCCTCCCTGGCCAATCTGTTT
 5 GAGGGGGAGGATGGCTCCCTTTCGCCCTCACC GGCTGATGCCAGTCGCCCTGCTGGCCCA
 GGCGATGGGCGACCAAATCTGCGCATGAAGTTCCAGGGCGCCTTCCGCAAGGGGGTGCCC
 AACCCCATCGATCTGCTGGAGTCCACCCTATATGAGTCCTCGGTGGTGCCTGGGCCCAAG
 AAAGCACCCATGGACTCACTGTTTGACTACGGCACCTATCGTCACCACTCCAGTGACAAC
 AAGAGGTGGAGGAAGAAGATCATAGAGAAGCAGCCGAGAGCCCCAAAGCCCCTGCCCCCT
 10 CAGCCGCCCCCATCCTCAAAGTCTTCAACCGGCCTATCCTCTTTGACATCGTGTCCTGG
 GGCTCCACTGCTGACCTGGACGGGCTGCTCCCATTTCTTGCTGACCCACAAGAAACGCCTA
 ACTGATGAGGAGTTTCGAGAGCCATCTACGGGGAAGACCTGCCTGCCCAAGGCCTTGCTG
 AACCTGAGCAATGGCCGCAACGACACCATCCCTGTGCTGCTGGACATCGCGGAGCGCACC
 GGCAACATGCGGGAGTTTCATTAACTCGCCCTTCCGTGACATCTACTATCGAGGTCAGACA
 15 GCCCTGCACATCGCCATTGAGCGTCGCTGCAAACACTACGTGGAACCTTCTCGTGCGCCAG
 GGAGCTGATGTCCACGCCCAGGCCCGTGGGCGCTTCTTCCAGCCCAAGGATGAGGGGGGC
 TACTTCTACTTTGGGGAGCTGCCCCGTGTCGCTGGCTGCCTGCACCAACCAGCCCCACATT
 GTCAACTACCTGACGGAGAACCCCCACAAGAAGGCGGACATGCGGCGCCAGGACTCGCGA
 GGCAACACAGTGCTGCATGCGCTGGTGGCCATTGCTGACAACACCCGTGAGAACACCAAG
 20 TTTGTTACCAAGATGTACGACCTGCTGCTGCTCAAGTGTGCCCCGCTCTTCCCCGACAGC
 AACCTGGAGGCCGTGCTCAACAACGACGGCCTCTCGCCCCCTCATGATGGCTGCCAAGACG
 GGCAAGATTGGGATCTTTTCAGCACATCATCCGGCGGGAGGTGACGGATGAGGACACACGG
 CACCTGTCCCGCAAGTTCAAGGACTGGGCCCTATGGGCCAGTGATTTCTCGCTTTATGAC
 CTCTCCTCCCTGGACACGTGTGGGGAAGAGGCCTCCGTGCTGGAGATCCTGGTGTACAAC
 25 AGCAAGATTGAGAACC GCCACGAGATGCTGGCTGTGGAGCCCATCAATGAACTGCTGCGG
 GACAAGTGGCGCAAGTTCCGGGGCCGTCTCCTTCTACATCAACGTGGTCTCCTACCTGTGT
 GCCATGGTCATCTTCACTCTCACCGCCTACTACCAGCCGCTGGAGGGCACACCGCCGTAC
 CCTTACCGCACACGGTGGACTACCTGCGGCTGGCTGGCGAGGTCAATTACGCTCTTCACT
 GGGGTCCTGTTCTTCTTTCACCAACATCAAAGACTTGTTCATGAAGAAATGCCCTGGAGTG
 30 AATTCTCTCTTCATTGATGGCTCCTTCCAGCTGCTCTACTTCATCTACTCTGTCTCTGGTG
 ATCGTCTCAGCAGCCCTCTACCTGGCAGGGATCGAGGCCTACCTGGCCGTGATGGTCTTT
 GCCCTGGTCTCTGGGCTGGATGAATGCCCTTTACTTTCACCCGTGGGCTGAAGCTGACGGGG
 ACCTATAGCATCATGATCCAGAAGATTCTCTTCAAGGACCTTTTCCGATTCTGTCTCGTC
 TACTTGCTCTTCATGATCGGCTACGCTTCAGCCCCTGGTCTCCCTCCTGAACCCGTGTGCC
 35 AACATGAAGGTGTGCAATGAGGACCAGACCAACTGCACAGTGCCCACTTACCCCTCGTGC
 CGTGACAGCGAGACCTTCAGCACCTTCCCTCCTGGACCTGTTTAAGCTGACCATTGGCATG
 GCGGACCTGGAGATGCTGAGCAGCACCAAGTACCCCGTGGTCTTCATCATCCTGTGGTG
 ACCTACATCATCCTCACCTTTGTGCTGCTCCTCAACATGCTCATTGCCCTCATGGGCGAG
 ACAGTGGGCCAGGTCTCCAAGGAGAGCAAGCACATCTGGAAGCTGCAGTGGGCCACCACC
 40 ATCCTGGACATTGAGCGCTCCTTCCCCGTATTCTCTGAGGAAGGCCTTCCGCTCTGGGGAG
 ATGGTCACCGTGGGCAAGAGCTCGGACGGCACCTCTGACCGCAGGTGGTGTCTCAGGGTG
 GATGAGGTGAACTGGTCTCACTGGAACCAGAACTTGGGCATCATCAACGAGGACCCGGGC

AAGAATGAGACCTACCAGTATTATGGCTTCTCGCATACCGTGGGCCGCTCCGCAGGGAT
CGCTGGTCCTCGGTGGTACCCCGCGTGGTGGAAGTGAACAAGAACTCGAACCCGGACGAG
GTGGTGGTGCCTCTGGACAGCATGGGGAACCCCGCTGCGATGGCCACCAGCAGGGTTAC
CCCCGCAAGTGGAGGACTGATGACGCCCCGCTCTAG

5

SEQ ID NO:2

MADSSEGPRAGPGEVAELPGDESGTPGGEAFPLSSLANLFEGEDGSLSPSPADASRPAGP
GDGRPNLRMKFQGAFRKGVNPIDLLESTLYESSVVPGPKKAPMDSLFDYGTyrRHSSDN
KRWRKKIIEKQPQSPKAPAPQPPILKVFNRPIFDIVSRGSTADLDGLLPFLLTHKKRL
10 TDEEFREPSTGKTCLPKALLNLSNGRNDTIPVLLDIAERTGNMREFINSPFRDIYYRGQT
ALHIAIERRCKHYVELLVAQGADVHAQARGRFFQPKDEGGYFYFGELPLSLAACTNQPHI
VNYLTENPHKKADMRRQDSRGNTVLHALVAIADNTRENTKFKVTKMYDLLLLKCARLFPDS
NLEAVLNNDGLSPLMMAAKTGKIGIFQHIIRREVTDEDTRHLSRKFKDWAYGPVYSSLYD
LSSLDTCGEEASVLEILVYNSKIENRHEMLAVEPINELLRDKWRKFGAVSFYINVVSYLE
15 AMVIFTLTAYYQPLEGTPPYPYRTTVDYLRRLAGEVITLFTGVLFFFTNIKDLFMKKCPGV
NSLFIDGSFQLLYFYISVLVIVSAALYLAGIEAYLAVMVFALVLGWMNALYFTRGLKLTG
TYSIMIQKILFKDLFRFLLVYLLFMIGYASALVSLNPNCANMKVCNEDQTNCTVPTYPSC
RDSETFSTFLLDLFLKTIGMGDLEMLSSTKYPVVFIIILLVTYIIILTFVLLLNMLIALMGE
TVGQVSKEKHIWKLQWATTILDIERSPVFLRKAFRSGEMVTVGKSSDGTDDRWCFRV
20 DEVNWSHWNQNLGIINEDPGKNETYQYYGFSHTVGRLRRDRWSSVPRVVELNKNNSNPDE
VVVPLDSMGNPRCDGHQQGYPRKWRTDDAPL

SEQ ID NO:3

CCACGCGTCCGCTCCCGGCCCGCGCGCCAGCCGTCCCGGAGGCTGAGCAGTGCAGACG
25 GGCCTGGGGCAGGCATGGCGGATTCCAGCGAAGGCCCGCGCGGGGCCCGGGAGGTGG
CTGAGCTCCCGGGGATGAGAGTGGCACCCAGGTGGGGAGGCTTTTCCTCTCTCCTCCC
TGGCCAATCTGTTTGAGGGGGGAGGATGGCTCCCTTTCGCCCTCACCGGCTGATGCCAGTC
GCCCTGCTGGCCCAGGCGATGGGCGACCAAATCTGCGCATGAAGTTCCAGGGCGCCTTCC
GCAAGGGGGTGCCCAACCCCATCGATCTGCTGGAGTCCACCCTATATGAGTCTCGGTGG
30 TGCCTGGGCCCCAAGAAAGCACCCATGGACTCACTGTTTGACTACGGCACCTATCGTCACC
ACTCCAGTGACAACAAGAGGTGGAGGAAGAAGATCATAGAGAAGCAGCCGAGAGCCCCA
AAGCCCCTGCCCTCAGCCGCCCCCATCTCAAAGTCTTCAACCGGCCTATCCTCTTTG
ACATCGTGTCCCGGGGCTCCACTGCTGACCTGGACGGGCTGCTCCCATCTTGCTGACCC
ACAAGAAACGCCTAACTGATGAGGAGTTTCGAGAGCCATCTACGGGGAAGACCTGCCTGC
35 CCAAGGCCTTGCTGAACCTGAGCAATGGCCGCAACGACACCATCCCTGTGCTGCTGGACA
TCGCGGAGCGCACCGGCAACATGCGGGAGTTTCATTAACTCGCCCTTCCGTGACATCTACT
ATCGAGGTGAGACAGCCCTGCACATCGCCATTGAGCGTCGCTGCAAACACTACGTGGAAC
TTCTCGTGGCCCAGGGAGCTGATGTCCACGCCAGGCCCGTGGGCGCTTCTTCCAGCCCCA
AGGATGAGGGGGGCTACTTCTACTTTGGGGAGCTGCCCCGTGCTGCTGGCTGCCTGCACCA
40 ACCAGCCCCACATTGTCAACTACCTGACGGAGAACCCCAAGAAGGCGGACATGCGGC
GCCAGGACTCGCGAGGCAACACAGTGCTGCATGCGCTGGTGGCCATTGCTGACAACACCC
GTGAGAACACCAAGTTTGTTACCAAGATGTACGACCTGCTGCTGCTCAAGTGTGCCCGCC

TCTTCCCCGACAGCAACCTGGAGGCCGTGCTCAACAACGACGGCCTCTCGCCCCCTCATGA
 TGGCTGCCAAGACGGGCAAGATTGGGATCTTTTCAGCACATCATCCGGCGGGAGGTGACGG
 ATGAGGACACACGGCACCTGTCCCGCAAGTTCAAGGACTGGGCCTATGGGCCAGTGATTT
 CCTCGCTTTATGACCTCTCCTCCCTGGACACGTGTGGGGAAGAGGCCTCCGTGCTGGAGA
 5 TCCTGGTGTTACAACAGCAAGATTGAGAACCGCCACGAGATGCTGGCTGTGGAGCCCATCA
 ATGAAGTGCTGCGGGACAAGTGGCGCAAGTTCGGGGCCGTCTCCTTCTACATCAACGTGG
 TCTCCTACCTGTGTGCCATGGTCATCTTCACTCTCACCGCCTACTACCAGCCGCTGGAGG
 GCACACCGCCGTACCCTTACCGCACCACGGTGGACTACCTGCGGCTGGCTGGCGAGGTCA
 TTACGCTCTTCACTGGGGTCTGTCTCTTCTTACCAACATCAAAGACTTGTTTCATGAAGA
 10 AATGCCCTGGAGTGAATTCTCTCTTCATTTGATGGCTCCTTCCAGCTGCTCTACTTTCATCT
 ACTCTGTCTCTGGTGATCGTCTCAGCAGCCCTCTACCTGGCAGGGATCGAGGCCTACCTGG
 CCGTGATGGTCTTTGCCCTGGTCTGGGCTGGATGAATGCCCTTTACTTTCACCCGTGGGC
 TGAAGCTGACGGGGACCTATAGCATCATGATCCAGAAGATTCTCTTCAAGGACCTTTTCC
 GATTCTGTCTCGTCTACTTGCTCTTCATGATCGGCTACGCTTCAGCCCTGGTCTCCCTCC
 15 TGAACCCGTGTGCCAACATGAAGGTGTGCAATGAGGACCAGACCAACTGCACAGTGCCCCA
 CTTACCCCTCGTGCCGTGACAGCGAGACCTTCAGCACCTTCCTCCTGGACCTGTTTAAGC
 TGACCATTGGCATGGGCGACCTGGAGATGCTGAGCAGCACCAAGTACCCCGTGGTCTTCA
 TCATCCTGCTGGTGACCTACATCATCCTCACCTTTGTGCTGCTCCTCAACATGCTCATTG
 CCCTCATGGGCGAGACAGTGGGCCAGGTCTCCAAGGAGAGCAAGCACATCTGGAAGCTGC
 20 AGTGGGCCACCACCATCCTGGACATTGAGCGCTCCTTCCCCGTATTCTTGAGGAAGGCCT
 TCCGCTCTGGGGAGATGGTCACCGTGGGCAAGAGCTCGGACGGCACTCCTGACCGCAGGT
 GGTGCTTCAGGGTGGATGAGGTGAACTGGTCTCACTGGAACCAGAACTTGGGCATCATCA
 ACGAGGACCCGGGCAAGAATGAGACCTACCAGTATTATGGCTTCTCGCATACCGTGGGCC
 GCCTCCGCAGGGATCGCTGGTCCCTCGGTGGTACCCCGCGTGGTGGAAGTGAACAAGAACT
 25 CGAACCCGGACGAGGTGGTGGTGCCTCTGGACAGCATGGGGAACCCCGCTGCGATGGCC
 ACCAGCAGGGTTACCCCGCAAGTGGAGGACTGATGACGCCCCGCTCTAGGGACTGCAGC
 CCAGCCCCAGCTTCTCTGCCCCACTCATTTCTAGTCCAGCCGCATTTTCAGCAGTGCCCTTCT
 GGGGTGTCCCCCACACCCTGCTTTGGCCCCAGAGGCGAGGGACCAGTGGAGGTGCCAGG
 GAGGCCCCAGGACCCTGTGGTCCCCCTGGCTCTGCCTCCCCACCCTGGGGTGGGGGCTCCC
 30 GGCCACCTGTCTTGCTCCTATGGAGTCACATAAGCCAACGCCAGAGCCCCCTCCACCTCAG
 GCCCCAGCCCCCTGCCTCTCCATTATTTATTTGCTCTGCTCTCAGGAAGCGACGTGACCCC
 TGCCCCAGCTGGAACCTGGCAGAGGCCTTAGGACCCCGTTCCAAGTGCACTGCCCCGCCA
 AGCCCCAGCCTCAGCCTGCGCCTGAGCTGCATGCGCCACCATTTTTTGGCAGCGTGGCAGC
 35 TTTTGCAAGGGGCTGGGGCCCTCGGCGTGGGGCCATGCCTTCTGTGTGTTCTGTAGTGTCT
 GGGATTTGCCGGTGCTCAATAAATGTTTATTCATTGACGGTGA

SEQ ID NO:4

CCGGCCGGGATTTCAGGAAGCGCGGATCTCCCGGCCCGCGGCCAGCCGTCCCGGAGGC
 TGAGCAGTGCAGACGGGCCTGGGGCAGGCATGGCGGATTCCAGCGAAGGCCCCCGCGCGG
 40 GGCCCGGGAGGTGGCTGAGCTCCCCGGGGATGAGAGTGGCACCCAGGTGGGGAGGCTT
 TTCTCTCTCCTCCCTGGCCAATCTGTTTGAGGGGGAGGATGGCTCCCTTTTCGCCCTCAC
 CGGCTGATGCCAGTCGCCCTGCTGGCCCAGGCGATGGGCGACCAAATCTGCGCATGAAGT

TCCAGGGCGCCTTCCGCAAGGGGGTGCCCAACCCCATCGATCTGCTGGAGTCCACCCTAT
 ATGAGTCCCTCGGTGGTGCCTGGGCCCCAAGAAAGCACCCATGGACTCACTGTTTGAAGTACG
 GCACCTATCGTCACCACTCCAGTGACAACAAGAGGTGGAGGAAGAAGATCATAGAGAAGC
 AGCCGCAGAGCCCCAAAGCCCCTGCCCCCTCAGCCGCCCCCATCCTCAAAGTCTTCAACC
 5 GGCCTATCCTCTTTGACATCGTGTCCCGGGGCTCCACTGCTGACCTGGACGGGCTGCTCC
 CATTCTTGCTGACCCACAAGAAACGCCTAACTGATGAGGAGTTTCGAGAGCCATCTACGG
 GGAAGACCTGCCTGCCCAAGGCCTTGCTGAACCTGAGCAATGGCCGCAACGACACCATCC
 CTGTGCTGCTGGACATCGCGGAGCGCACCGGCAACATGCGGGAGTTCAATTAACCTCGCCCT
 TCCGTGACATCTACTATCGAGGTCAGACAGCCCTGCACATCGCCATTGAGCGTCGCTGCA
 10 AACACTACGTGGAACCTTCTCGTGGCCAGGGAGCTGATGTCCACGCCCAGGCCCGTGGGC
 GCTTCTTCCAGCCCCAAGGATGAGGGGGGCTACTTCTACTTTGGGGAGCTGCCCCCTGTCGC
 TGGCTGCCTGCACCAACCAGCCCCACATTGTCAACTACCTGACGGAGAACCCCCACAAGA
 AGGCGGACATGCGGCGCCAGGACTCGCGAGGCAACACAGTGCTGCATGCGCTGGTGGCCA
 TTGCTGACAACACCCGTGAGAACACCAAGTTTGTACCAAGATGTACGACCTGCTGCTGC
 15 TCAAGTGTGCCCCGCTCTTCCCCGACAGCAACCTGGAGGCCGTGCTCAACAACGACGGCC
 TCTCGCCCCCTCATGATGGCTGCCAAGACGGGCAAGATTGGGATCTTTCAGCACATCATCC
 GCGGGAGGTGACGGATGAGGACACACGGCACCTGTCCCGCAAGTTCAAGGACTGGGCCT
 ATGGGCCAGTGATTTCTCGCTTTATGACCTCTCCTCCCTGGACACGTGTGGGGAAGAGG
 CCTCCGTGCTGGAGATCCTGGTGTACAACAGCAAGATTGAGAACCGCCACGAGATGCTGG
 20 CTGTGGAGCCCATCAATGAAGTGTGCGGGACAAGTGGCGCAAGTTGCGGGCCGTCTCCT
 TCTACATCAACGTGGTCTCCTACCTGTGTGCCATGGTCATCTTCACTCTCACCGCCTACT
 ACCAGCCGCTGGAGGGCACACCGCCGTACCCTTACCGCACCACGGTGGACTACCTGCGGC
 TGGCTGGCGAGGTCAATTACGCTCTTCACTGGGGTCCTGTTCTTCTTACCAACATCAAAG
 ACTTGTTTCATGAAGAAATGCCCTGGAGTGAATTCCTCTCTTCAATTGATGGCTCCTTCCAGC
 25 TGCTCTACTTCATCTACTCTGTCTTGGTGATCGTCTCAGCAGCCCTCTACCTGGCAGGGA
 TCGAGGCCTACCTGGCCGTGATGGTCTTTGCCCCGGTCCCTGGGCTGGATGAATGCCCTTT
 ACTTCACCCGTGGGCTGAAGCTGACGGGGACCTATAGCATCATGATCCAGAAGATTCTCT
 TCAAGGACCTTTTCCGATTCTGCTCGTCTACTTGCTCTTCATGATCGGCTACGCTTCAG
 CCCTGGTCTCCCTCCTGAACCCGTGTGCCAACATGAAGGTGTGCAATGAGGACCAGACCA
 30 ACTGCACAGTGCCCACTTACCCCTCGTGCCGTGACAGCGAGACCTTCAGCACCTTCTCTCC
 TGGACCTGTTTAAAGCTGACCATTTGGCATGGGCGACCTGGAGATGCTGAGCAGCACCAAGT
 ACCCCGTGGTCTTCATCATCCTGCTGGTGACCTACATCATCCTCACCTTTGTGCTGCTCC
 TCAACATGCTCATTTGCCCTCATGGGCGAGACAGTGGGCCAGGTCTCCAAGGAGAGCAAGC
 ACATCTGGAAGCTGCAGTGGGCCACCACCATCCTGGACATTGAGCGCTCCTTCCCCGTAT
 35 TCCTGAGGAAGGCCTTCCGCTCTGGGGAGATGGTACCGTGGGCAAGAGCTCGGACGGCA
 CTCCTGACCGCAGGTGGTGTCTCAGGTGGATGAGGTGAACTGGTCTCACTGGAACCAGA
 ACTTGGGCATCATCAACGAGGACCCGGGCAAGAATGAGACCTACCAGTATTATGGCTTCT
 CGCATACCGTGGGCCCGCTCCGCAGGGATCGCTGGTCTCCTCGGTGGTACCCCGCGTGGTG
 AACTGAACAAGAACTCGAACCCGGACGAGGTGGTGGTGCCTCTGGACAGCATGGGGAACC
 40 CCCGCTGCGATGGCCACCAGCAGGGTTACCCCGCAAGTGGAGGACTGATGACGCCCCGC
 TCTAGGACTGCAGCCAGCCCCAGCTTCTCTGCCACTCATTTCTAGTCCAGCCGCATT
 TCAGCAGTGCTTCTGGGGTGTCCCCCACACCCTGCTTTGGCCCCAGAGGCGAGGGACC

AGTGGAGGTGCCAGGGAGGCCCCAGGACCCTGTGGTCCCCTGGCTCTGCCTCCCCACCCT
GGGGTGGGGGCTCCCGGCCACCTGTCTTGCTCCTATGGAGTCACATAAGCCAACGCCAGA
GCCCCCTCCACCTCAGGCCCCAGCCCCTGCTCTCCATTATTTATTTGCTCTGCTCTCAGG
AAGCGACGTGACCCCTGCCCCAGCTGGAACCTGGCAGAGGCCTTAGGACCCCGTTCCAAG
5 TGCAC TGCCCGGCCAAGCCCCAGCCTCAGCCTGCGCCTGAGCTGCATGCGCCACCATTTT
TGGCAGCGTGGCAGCTTTGCAAGGGGCTGGGGCCCTCGGCGTGGGGCCATGCCTTCTGTG
TGTTCGTAGTGTCTGGGATTTGCCGGTGCTCAATAAATGTTTATTCATTGACGGTG